

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/236967373>

# Sugar addiction: Pushing the drug-sugar analogy to the limit

ARTICLE · MAY 2013

DOI: 10.1097/MCO.0b013e328361c8b8 · Source: PubMed

---

CITATIONS

27

---

READS

2,394

## 3 AUTHORS:



[Serge H Ahmed](#)

French National Centre for Scientific Resea...

86 PUBLICATIONS 3,862 CITATIONS

SEE PROFILE



[Karine Guillem](#)

Université Victor Segalen Bordeaux 2

33 PUBLICATIONS 427 CITATIONS

SEE PROFILE



[Youna Vandaele](#)

Johns Hopkins University

3 PUBLICATIONS 28 CITATIONS

SEE PROFILE



# Sugar addiction: pushing the drug-sugar analogy to the limit

Serge H. Ahmed<sup>a,b</sup>, Karine Guillem<sup>a,b</sup>, and Youna Vandaele<sup>a,b</sup>

## Purpose of review

To review research that tests the validity of the analogy between addictive drugs, like cocaine, and hyperpalatable foods, notably those high in added sugar (i.e., sucrose).

## Recent findings

Available evidence in humans shows that sugar and sweetness can induce reward and craving that are comparable in magnitude to those induced by addictive drugs. Although this evidence is limited by the inherent difficulty of comparing different types of rewards and psychological experiences in humans, it is nevertheless supported by recent experimental research on sugar and sweet reward in laboratory rats. Overall, this research has revealed that sugar and sweet reward can not only substitute to addictive drugs, like cocaine, but can even be more rewarding and attractive. At the neurobiological level, the neural substrates of sugar and sweet reward appear to be more robust than those of cocaine (i.e., more resistant to functional failures), possibly reflecting past selective evolutionary pressures for seeking and taking foods high in sugar and calories.

## Summary

The biological robustness in the neural substrates of sugar and sweet reward may be sufficient to explain why many people can have difficulty to control the consumption of foods high in sugar when continuously exposed to them.

## Keywords

addiction, animal models, cocaine, craving, dopamine, sugar

## INTRODUCTION

The current global increase in obesity prevalence and the difficulty of containing it despite the negative consequences have recently led several researchers, mostly neuroscientists, to compare obesity to drug addiction [1–9,10<sup>■</sup>,11] and palatable foods, particularly those high in added sugar (i.e., sucrose), to addictive drugs like cocaine [4,12–14]. Volkow – the current head of the American National Institute on Drug Abuse – and O’Brien [9] were among the first to suggest that the concept of addiction may shed some new light on obesity – an idea that has proven quite influential, as shown by the exponential rise of the use of the expression ‘food addiction’ (and related terms) in the biomedical and scientific literature ever since [15<sup>■</sup>]. In parallel, others have strongly argued that hyperpalatable foods rich in added sugar and/or fat could be genuinely addictive, at least in a significant proportion of exposed people [12,13]. One currently estimates that about 10–20% of people would present addiction-like symptoms

toward hyperpalatable foods [16,17] – a proportion that is not different from the proportion of cocaine or heroin users who go on to develop addiction [18]. The widespread introduction of hyperpalatable foods during the 20th century could be likened to the introduction of distilled drinks (i.e., gins, whiskeys) in the 17th century or of injectable synthetic drugs at the end of the 19th century, each spurred its own addiction epidemics [19]. Finally, people are as ill prepared biologically to foods high in added sugar and/or fat, as they are to drugs in pure or highly concentrated form [4]. In this regard, the ubiquity, ready availability, and affordability of

<sup>a</sup>Université de Bordeaux and <sup>b</sup>CNRS, Institut des Maladies Neurodégénératives, Bordeaux, France

Correspondence to Serge H. Ahmed, Université de Bordeaux, Institut des Maladies Neurodégénératives, UMR 5293, 146 rue Léo-Saignat, F-33000 Bordeaux, France. Tel: +33 557 571 566; fax: +33 556 900 278; e-mail: sahmed@u-bordeaux2.fr

**Curr Opin Clin Nutr Metab Care** 2013, 16:434–439

DOI:10.1097/MCO.0b013e328361c8b8

## KEY POINTS

- Sugar and sweetness can induce reward and craving in humans that are at least comparable in magnitude to those induced by addictive drugs.
- Sugar and sweet reward cannot only substitute to cocaine but can even be more rewarding and attractive than cocaine in animals.
- The neural substrates of sugar and sweet reward are more robust than those of cocaine (i.e., more resistant to functional failures) in animals.
- Additional research using more valid psychophysical and behavioral methods is required in humans to more directly compare addictive drugs and sugar.

those foods make them a serious modern hazard to public health [1,10<sup>¶</sup>].

However, although the concept of food and sugar addiction is gaining momentum, it is also currently a subject of intense debate and no solid evidence-based consensus has emerged yet [20–22,23<sup>¶</sup>,24<sup>¶¶</sup>]. What is really at stake is, first, the drug-like status of some food ingredients, notably sugar, and, second, the relevance of the concept of addiction to understanding obesity. The latter issue has been recently discussed at length elsewhere, particularly the overlap in the neurobiological substrates between drug addiction and obesity [10<sup>¶</sup>,12,13,23<sup>¶</sup>]. Here we will be mainly concerned with the strengths and limitations of the analogy between drugs of abuse and hyperpalatable foods, with a particular, although not exclusive, focus on foods or drinks containing high levels of added sugar (i.e., sucrose). The world history of sugar is not different from that of many psychoactive drugs, including cocaine. It initially began as a medicine for the rich and the powerful and ended up as a product of mass consumption [19]. Today, the ‘sweetening of the world’s diet’ is almost total [25] and there is growing evidence linking increased sugar availability and consumption to overweight and obesity [26,27].

## DRUG VERSUS FOOD PSYCHOACTIVE INTOXICATION

At first glance, the analogy between foods high in added sugar and drugs of abuse, such as cocaine, may seem overstretched, not to say absurd. Unlike sugar, drugs of abuse, except ethanol, are non-nutritive molecules that, once self-administered (e.g., through inhalation or via the intravenous route), quickly cross the blood-brain barrier to

physically interact and interfere with specific endogenous molecular substrates and processes, generally at the surface of brain cells [28]. For instance, cocaine binds to the dopamine transporter, among other molecular targets, thereby blocking the neuronal reuptake of dopamine and causing an abnormally high surge of dopamine in innervated brain regions [29]. Drug-induced changes in neuronal and synaptic activity in different brain regions and circuits lead in turn to alterations in behavioral dispositions (e.g., aggressiveness, risk-taking), mood (e.g., euphoria) and other mental functions (e.g., judgement, decision-making) [30]. At high doses, these psychoactive effects may considerably impair normal functions (e.g., distorted perception; altered judgement; diminished self-control) and can be hazardous to both the self and others. In contrast, other psychoactive effects can be advantageous or functional under some circumstances (e.g., disinhibition of sex) and, of course, even highly stimulating and rewarding (e.g., euphoria) [31].

Foods high in sugar can also change brain activity but via more natural routes than drugs of abuse. They can change brain activity, first, via the stimulation of specialized sweet taste cells in the mouth (and in the gut) [32,33<sup>¶¶</sup>] and, second, via postabsorptive brain mechanisms involving glucose signaling [34] – the latter being the most drug-like. Nevertheless, people now often report seeking and consuming sweet foods for their drug-like psychoactive and mood-altering effects [35]. They eat sweet foods to experience highly rewarding sensations, to cope with stress (e.g., stress or comfort eating), pain or fatigue, to enhance cognition and/or to ameliorate bad mood (e.g., relief of negative affect). This anecdotal evidence is also confirmed by research in different populations (e.g., adolescents at risk to develop depression; obese women) showing that sweet foods can indeed elicit different desirable drug-like psychoactive effects, including affective comfort and alleviation of depressed mood [35,36–38]. In most cases, however, the magnitude of the experienced psychoactive effects of sweet foods is mild and does not seem to match those of drugs. In other words, sweet foods are clearly not as behaviorally and/or psychologically toxic as drugs of abuse can be, especially at high doses. For instance, unlike drugs, consumption of hyperpalatable foods, even extremely high in sugar, does not produce any abnormal mental state or change in behavioral disposition. This explains why no policeman will arrest a driver because he/she had eaten several doughnuts before driving his/her car or why no judge will consider drinking half a gallon of sugar-sweetened soda before committing a crime

as a mitigating circumstance. If any, foods rich in sugar seem to produce more advantageous effects on decision-making and self-control than disadvantageous ones. For instance, sugar has been shown to boost self-control under some circumstances [39]. Sugar and sweetness can also promote helping attitude and thereby encourage cooperation among people [40].

### **DRUG VERSUS FOOD REWARD**

The analogy between foods high in sugar and drugs of abuse, like cocaine, may also seem exaggerated because drug reward is generally thought to be incommensurably more intense than food reward. This difference is generally attributed to drugs' ability to activate brain reward circuits, notably midbrain dopamine neurons, more potently than any other nondrug reward [10<sup>9</sup>,41]. Drug addicts often report that the first drug experiences can be even better than sex orgasm. In Western societies, sex orgasm is generally placed at the top of the human hedonic scale, well above foods [42]. This is memorably illustrated in the movie 'Meet Joe Black' where Joe Black – who has recently developed a taste and craving for peanut butter – confesses to Suzan Parrish that making love with her was better than peanut butter. In fact, however, there is little direct evidence showing that drug reward is indeed more intense than food or sex reward, even in drug addicts. In one unique comparative study that has begun to address this question in cocaine addicts, addicts reported liking food as much as they liked cocaine or sex [43]. Interestingly, in this study, healthy individuals also reported liking food almost equally as sex. More research is clearly needed here to address this important issue about the relative intensity of food versus drug reward (see also below).

### **DRUG VERSUS FOOD CRAVING**

The analogy between hyperpalatable foods high in sugar and drugs of abuse, like cocaine, may also seem absurd because drug craving is by definition a pathological desire for a specific substance that is unwanted but difficult to resist [44]. Can food craving be as intense as drug craving? Once again there is little empirical comparative research on this question. Most people do report experiencing food cravings, sometimes very intense ones [45]. Chocolate and sweet foods are the most common craved foods [35,46,47]. In a recent large-scale experience sampling study on everyday desires and urges, food desires were by far the most frequent, with a large proportion of these desires being felt as conflictual and eliciting resistance

[48<sup>9</sup>]. The key issue here, however, is to determine whether food cravings can be comparable in magnitude to drug cravings. In one rare study addressing this problem, cocaine addicts reported wanting food as intensely as they wanted cocaine or sex [43]. In another recent study, craving for a cigarette or for a palatable food was induced in chronic smokers by exposure to cigarette or food cues, respectively. Overall, the intensity and resistibility (i.e., ability to resist it) of food craving were comparable to those of cigarette craving [49]. At the neurobiological level, a recent meta-analysis of neuroimaging research revealed a large overlap in the brain networks underlying cue-induced food versus cigarette craving [50<sup>99</sup>]. A similar overlap in neural substrates is also observed in animal models of food and drug craving [51].

### **WHEN SUGAR AND SWEET REWARD SURPASSES COCAINE REWARD**

Overall, available research shows that, although hyperpalatable foods high in sugar are clearly not as behaviorally and psychologically toxic as cocaine and other drugs of abuse, sweet reward and craving are apparently comparable in intensity or magnitude to drug reward and craving, thereby providing some support to the analogy between hyperpalatable foods and drugs. However, solid evidence for the food-drug analogy is still scant and most of it is based on poorly validated intersubjective comparisons and evaluations by people with drug addiction who are clearly not representative of the general population currently exposed to foods high in sugar. There are valid psychophysical methods that could allow one to compare drug reward (or craving) with food reward (or craving) but these methods have yet to be tested [52]. Similarly, there are objective behavioral methods for measuring reinforcing strengths between different kinds of rewards but they have not been systematically used to directly compare hyperpalatable foods with drugs of abuse [53]. Finally, at the neurobiological level, drug and food cues clearly recruit the same overall brain networks [50<sup>99</sup>] but more direct quantitative comparisons between brain activation patterns during drug versus food reward (or craving) are still lacking.

Another possible approach to explore the drug-food analogy could be to compare drug versus food reward in nonhuman animals, such as laboratory rats or mice. Rats have been shown to self-administer most addictive drugs, including cocaine [54], and to develop most of the behavioral signs of addiction after prolonged drug self-administration [55]. Like humans, rats have also an inborn sweet

tooth that has been systematically exploited over the past 60 years for research on the neural basis of reward and motivation. In fact, many important discoveries about the neurobiology of reward and motivation have been and are still made using sugar and sweetness as a reward or incentive [2]. Sugar and sweetness are also often used in drug addiction research to train animals before drug self-administration training. More relevant to the present review, under certain circumstances, sugar and sweet reward can substitute to cocaine, thereby decreasing cocaine self-administration [11]. Furthermore, when directly compared together, sugar and sweet reward can even be more rewarding and attractive than cocaine [14,56–59]. For instance, in a recent series of choice experiments, we found that when rats are offered an exclusive choice between sucrose (or saccharin) and cocaine self-administration, they develop a strong and persistent preference for sucrose (or saccharin) [14,56–58]. Similar findings have also been obtained in rats offered an exclusive choice between nicotine and sucrose [60]. Sugar withdrawal can also induce behavioral and neurochemical signs similar to those of heroin withdrawal [12]. Table 1 summarizes behavioral evidence from different studies that have compared sucrose (or sugar-sweetened foods) with cocaine under similar testing conditions. Clearly, rats can be at least as rewarded or motivated for sucrose as for cocaine and sometimes even more.

At first glance, this conclusion seems to conflict with evidence showing that sugar and sweet reward are much less potent than cocaine to boost brain dopamine signaling [14]. However, this apparent discrepancy may also suggest that dopamine is probably not enough to drive preference and that sugar, unlike cocaine reward, involves more than brain dopamine [14,24<sup>11</sup>]. This interpretation is supported by recent research using optogenetic methods in mice. Mice were allowed to choose between two sippers: licking one sipper delivered water and optogenetic stimulation of dopaminergic neurons whereas licking the other sipper delivered water sweetened with sucrose. When concentrations were sufficiently high, mice preferred sucrose over optogenetic stimulation of dopamine neurons [61] which can be rewarding alone [62]. In addition, there is ample, although disparate and overlooked, evidence that the neurobiological substrate of sucrose reward is more robust than that of cocaine reward. Table 2 summarizes the outcomes of all published studies ( $n=75$ ) that have compared the effects of different genetic (e.g., gene knockout or knockdown), pharmacological (e.g., selective receptor antagonism) or neurobiological (e.g., selective brain lesion) interventions on cocaine versus

**Table 1. Sucrose reward and motivation versus cocaine reward and motivation**

Behavioral criterion	Sucrose/Sweet	Cocaine
Preference	++	+
Demand inelasticity	++	+
Punishment resistance	+	+
Breaking point	++	++

Preference refers to sucrose (or sweet) choice over cocaine in choice experiments [14,56–58]. Demand inelasticity refers to the degree of resistance of sucrose or cocaine consumption to increasing behavioral costs (e.g., increasing number of required responses) [63–65]. Similarly, resistance to punishment refers to the degree of resistance of sucrose or cocaine consumption to aversive electrical footshock or associated conditioned stimuli [66–70]. Finally, breaking point refers to the maximum acceptable work before giving up working for sucrose (or sweet reward) or cocaine [57].

sucrose under similar behavioral conditions (e.g., fixed-ratio or progressive-ratio schedule of reinforcement). Overall, most interventions affected cocaine-rewarded or cocaine-motivated behaviors (i.e., 88 out of a total of 91 interventions) but only a few of them (i.e., 17) impacted on sucrose-rewarded or

**Table 2. Neurobiological interventions on sucrose versus cocaine reward or motivation**

Interventions	<i>N</i>	Sucrose	Cocaine
Intracellular processes	8	25	100
Selective brain lesions	8	12.5	87.5
Neurotransmitters			
Dopamine	23	21.7	100
Glutamate	21	14.3	90
Serotonin	8	25	100
Acetylcholine	6	33.3	100
Neuropeptides	6	16.6	100
Adenosine	3	0	100
Norepinephrine	2	0	100
Cannabinoids	2	0	100
Steroids	4	25	100
Total	91	18.7	96.7

We conducted a search for articles containing the words ‘sucrose’ and ‘cocaine’ in the PubMed database (last accession date: March 11, 2013). A total of 256 articles were retrieved. Among these 256 articles, 75 were selected and retained as relevant to the present study based on information contained in the abstract. The full-text of the selected articles was then read and analyzed in-depth for interventions that were tested on both sucrose and cocaine reward (or motivation) and under similar experimental conditions (e.g., intervention X tested on both operant responding for sucrose and cocaine under a comparable schedule of reinforcement). The column *N* shows the number of interventions *per* specific biological target. The number in the other columns indicates the proportion of interventions in % that affected sucrose- or cocaine-related behaviors. Note that some articles can contain more than one neurobiological intervention. The list of selected articles is available upon request to the authors.

sucrose-motivated behaviors. When the effect of each specific intervention is considered individually, it is generally interpreted as evidence that a given gene, brain region or neurotransmitter system is specifically involved in cocaine reward or motivation but not in sucrose reward or motivation. However, the big picture emerging from Table 2 clearly shows that many different specific interventions can profoundly affect cocaine reward or motivation while sparing sucrose reward or motivation. This discovery strongly suggests that the neurobiological substrate of sugar reward and motivation is more robust than that of cocaine (i.e., more resistance to individual biological failures or dysfunctions). Such robustness probably results from the evolution of multiple neural mechanisms for seeking and taking sugar and other foods high in calories [24<sup>■</sup>] and further underscores the exceptional motivational power of sugar sweetness [14].

## CONCLUSION

The extreme intensity of reward and craving produced by addictive drugs, like cocaine, largely explains, together with their toxic effects, why these drugs tend to be universally prohibited. The danger to be avoided by prohibition is that drug reward will turn people away from more socially valued, healthy, and productive life activities and occupations. Most of the current medical diagnostic criteria of drug addictive disorders reflect this harmful shift in behavioral preference and priority [44]. As reviewed here, there is now strong evidence supporting the notion that hyperpalatable foods, notably those high in added sugar, can induce reward and craving that are at least comparable to addictive drugs. Although more research in humans is clearly needed to confirm this conclusion, there is now solid evidence in nonhuman animals showing that sugar and sweet reward can even be more rewarding and attractive than addictive drugs, probably owing to an underlying robust neural substrate. Such biological robustness may be sufficient to explain why people can have difficulty to control the consumption of foods high in sugar when continuously exposed to them.

## Acknowledgements

*This work is supported by the French Research Council (CNRS), the French National Agency (ANR), the Fondation NRJ, the Université Bordeaux-Segalen and the Conseil Regional d'Aquitaine.*

## Conflicts of interest

*There are no conflicts of interest.*

## REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

Additional references related to this topic can also be found in the Current World Literature section in this issue (p. 492).

1. Allen PJ, Batra P, Geiger BM, *et al.* Rationale and consequences of reclassifying obesity as an addictive disorder: neurobiology, food environment and social policy perspectives. *Physiol Behav* 2012; 107:126–137.
2. Berridge KC, Ho CY, Richard JM, DiFeliceantonio AG. The tempted brain eats: pleasure and desire circuits in obesity and eating disorders. *Brain Res* 2010; 1350:43–64.
3. Davis C, Curtis C, Levitan RD, *et al.* Evidence that 'food addiction' is a valid phenotype of obesity. *Appetite* 2011; 57:711–717.
4. Ifland JR, Preuss HG, Marcus MT, *et al.* Refined food addiction: a classic substance use disorder. *Med Hypotheses* 2009; 72:518–526.
5. Kenny PJ. Reward mechanisms in obesity: new insights and future directions. *Neuron* 2011; 69:664–679.
6. Liu Y, von Deneen KM, Kobeissy FH, Gold MS. Food addiction and obesity: evidence from bench to bedside. *J Psychoactive Drugs* 2011; 42:133–145.
7. Parylak SL, Koob GF, Zorrilla EP. The dark side of food addiction. *Physiol Behav* 2011; 104:149–156.
8. Pelchat ML. Food addiction in humans. *J Nutr* 2009; 139:620–622.
9. Volkow ND, O'Brien CP. Issues for DSM-V: should obesity be included as a brain disorder? *Am J Psychiatry* 2007; 164:708–710.
10. Volkow ND, Wang GJ, Tomasi D, Baler RD. The Addictive Dimensionality of ■ Obesity. *Biol Psychiatry* 2013. (in press).  
An up-to-date review of the overlap in neurobiological substrates between drug addiction and obesity.
11. Ahmed SH, Avena NM, Berridge KC, Gearhardt AN, Guillem K. Food addiction. In *Neuroscience in the 21st Century: from basic to clinical*. Edited by Pfaff DW. New York: Springer; 2012:2833–2857.
12. Avena NM, Rada P, Hoebel BG. Evidence for sugar addiction: behavioral and neurochemical effects of intermittent, excessive sugar intake. *Neurosci Biobehav Rev* 2008; 32:20–39.
13. Gearhardt AN, Grilo CM, Dileone RJ, *et al.* Can food be addictive? Public health and policy implications. *Addiction* 2011; 106:1208–1212.
14. Lenoir M, Serre F, Cantin L, Ahmed SH. Intense sweetness surpasses cocaine reward. *PLoS One* 2007; 2:e698.
15. Salamone JD, Correa M. Dopamine and Food Addiction: Lexicon Badly ■ Needed. *Biol Psychiatry* 2013. (in press).  
A healthy reminder that words like 'reward' and 'addiction' have specific technical meanings and need to be used with great caution and moderation to avoid confusion.
16. Gearhardt AN, Corbin WR, Brownell KD. Preliminary validation of the Yale Food Addiction Scale. *Appetite* 2009; 52:430–436.
17. Meule A. How Prevalent is 'Food Addiction'? *Front Psychiatry* 2011; 2:61.
18. Sussman S, Lisha N, Griffiths M. Prevalence of the addictions: a problem of the majority or the minority? *Eval Health Prof* 2011; 34:3–56.
19. Courtwright DT. Addiction and the science of history. *Addiction* 2012; 107:486–492.
20. Benton D. The plausibility of sugar addiction and its role in obesity and eating disorders. *Clin Nutr* 2010; 29:288–303.
21. Corsica JA, Pelchat ML. Food addiction: true or false? *Curr Opin Gastroenterol* 2011; 26:165–169.
22. Wilson GT. Eating disorders, obesity and addiction. *Eur Eat Disord Rev* 2010; 18:341–351.
23. Ziauddeen H, Farooqi IS, Fletcher PC. Obesity and the brain: how convincing ■ is the addiction model? *Nat Rev Neurosci* 2012; 13:279–286.  
This critical review argues that the addiction model of obesity is not yet backed by solid scientific evidence and seriously questions the validity and relevance of current animal models of food addiction to the human condition.
24. DiLeone RJ, Taylor JR, Picciotto MR. The drive to eat: comparisons ■ and distinctions between mechanisms of food reward and drug addiction. *Nat Neurosci* 2012; 15:1330–1335.  
This review provides a recent analysis of both behavioral and neurobiological commonalities and differences between drugs of abuse and palatable foods (e.g., rich in sugar). Notably, the authors suggest that selective pressures may have led to the evolution of multiple neural mechanisms for seeking and taking sweet and highly caloric foods.
25. Popkin BM, Adair LS, Ng SW. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev* 2012; 70:3–21.
26. Malik VS, Hu FB. Sweeteners and risk of obesity and type 2 diabetes: the role of sugar-sweetened beverages. *Curr Diab Rep* 2012; 12:195–203.
27. de Ruyter JC, Olthof MR, Seidell JC, Katan MB. A trial of sugar-free or sugar-sweetened beverages and body weight in children. *N Engl J Med* 2012; 367:1397–1406.
28. Hanson GR, Venturelli PJ, Fleckenstein AE. *Drugs and society*. 9th edn Boston: Jones and Bartlett; 2006. p. 575.

29. Sulzer D. How addictive drugs disrupt presynaptic dopamine neurotransmission. *Neuron* 2011; 69:628–649.
30. Volkow ND, Baler RD, Goldstein RZ. Addiction: pulling at the neural threads of social behaviors. *Neuron* 2011; 69:599–602.
31. Müller CP, Schumann G. Drugs as instruments: a new framework for nonaddictive psychoactive drug use. *Behav Brain Sci* 2011; 34:293–310.
32. Yarmolinsky DA, Zuker CS, Ryba NJ. Common sense about taste: from mammals to insects. *Cell* 2009; 139:234–244.
33. Brown RJ, Rother KI. Nonnutritive sweeteners and their role in the gastrointestinal tract. *J Clin Endocrinol Metab* 2012; 97:2597–2605.
- An excellent overview of the biology of intestinal sweet taste receptors and of their role in modulating glucose absorption.
34. Grayson BE, Seeley RJ, Sandoval DA. Wired on sugar: the role of the CNS in the regulation of glucose homeostasis. *Nat Rev Neurosci* 2013; 14:24–37.
35. Pretlow RA. Addiction to highly pleasurable food as a cause of the childhood obesity epidemic: a qualitative Internet study. *Eat Disord* 2011; 19:295–307.
36. Rose N, Koperski S, Golomb BA. Mood food: chocolate and depressive symptoms in a cross-sectional analysis. *Arch Intern Med* 2010; 170:699–703.
37. Dallman MF. Stress-induced obesity and the emotional nervous system. *Trends Endocrinol Metab* 2011; 21:159–165.
38. Spring B, Schneider K, Smith M, *et al.* Abuse potential of carbohydrates for overweight carbohydrate cravers. *Psychopharmacology (Berl)* 2008; 197:637–647.
39. Gailliot MT, Baumeister RF, DeWall CN, *et al.* Self-control relies on glucose as a limited energy source: willpower is more than a metaphor. *J Pers Soc Psychol* 2007; 92:325–336.
40. Meier BP, Moeller SK, Riemer-Peltz M, Robinson MD. Sweet taste preferences and experiences predict prosocial inferences, personalities, and behaviors. *J Pers Soc Psychol* 2012; 102:163–174.
41. Smith DG, Robbins TW. The neurobiological underpinnings of obesity and binge eating: a rationale for adopting the food addiction model. *Biol Psychiatry* 2013. (in press).
42. Kenrick DT, Griskevicius V, Neuberg SL, Schaller M. Renovating the pyramid of needs: contemporary extensions built upon ancient foundations. *Perspect Psychol Sci* 2010; 5:292–314.
43. Goldstein RZ, Woicik PA, Moeller SJ, *et al.* Liking and wanting of drug and nondrug rewards in active cocaine users: the STRAP-R questionnaire. *J Psychopharmacol* 2010; 24:257–266.
44. DSM-5. The Future of Psychiatric Diagnosis on World Wide Web URL: <http://www.dsm5.org/Pages/Default.aspx>. [Accessed 9 March 2013]
45. Hill AJ. The psychology of food craving. *Proc Nutr Soc* 2007; 66:277–285.
46. Meule A, Kubler A. Food cravings in food addiction: the distinct role of positive reinforcement. *Eat Behav* 2012; 13:252–255.
47. Davis C, Carter JC. Compulsive overeating as an addiction disorder. A review of theory and evidence. *Appetite* 2009; 53:1–8.
48. Hofmann W, Baumeister RF, Forster G, Vohs KD. Everyday temptations: an experience sampling study of desire, conflict, and self-control. *J Pers Soc Psychol* 2012; 102:1318–1335.
- This interesting study used the mobile smartphone technology to follow and record everyday temptations in about 200 people over a 1-week period.
49. Kober H, Mende-Siedlecki P, Kross EF, *et al.* Prefrontal-striatal pathway underlies cognitive regulation of craving. *Proc Natl Acad Sci U S A* 2010; 107:14811–14816.
50. Tang DW, Fellows LK, Small DM, Dagher A. Food and drug cues activate similar brain regions: a meta-analysis of functional MRI studies. *Physiol Behav* 2012; 106:317–324.
- This important meta-analysis of human neuroimaging studies reveals a large overlap in neural substrates between food cravings and drug cravings (i.e., cigarette smoking).
51. Nair SG, Adams-Deutsch T, Epstein DH, Shaham Y. The neuropharmacology of relapse to food seeking: methodology, main findings, and comparison with relapse to drug seeking. *Prog Neurobiol* 2009; 89:18–45.
52. Lim J, Wood A, Green BG. Derivation and evaluation of a labeled hedonic scale. *Chem Senses* 2009; 34:739–751.
53. Epstein LH, Leddy JJ, Temple JL, Faith MS. Food reinforcement and eating: a multilevel analysis. *Psychol Bull* 2007; 133:884–906.
54. O'Connor EC, Chapman K, Butler P, Mead AN. The predictive validity of the rat self-administration model for abuse liability. *Neurosci Biobehav Rev* 2010; 35:912–938.
55. Ahmed SH. The science of making drug-addicted animals. *Neuroscience* 2012; 211:107–125.
56. Augier E, Vouillac C, Ahmed SH. Diazepam promotes choice of abstinence in cocaine self-administering rats. *Addict Biol* 2012; 17:378–391.
57. Cantin L, Lenoir M, Augier E, *et al.* Cocaine is low on the value ladder of rats: possible evidence for resilience to addiction. *PLoS One* 2010; 5:e11592.
58. Kerstetter KA, Ballis MA, Duffin-Lutgen S, *et al.* Sex differences in selecting between food and cocaine reinforcement are mediated by estrogen. *Neuropsychopharmacology* 2012; 37:2605–2614.
59. Morales L, Del Olmo N, Valladolid-Acebes I, *et al.* Shift of circadian feeding pattern by high-fat diets is coincident with reward deficits in obese mice. *PLoS One* 2012; 7:e36139.
60. Le Sage MG. Toward a nonhuman model of contingency management: effects of reinforcing abstinence from nicotine self-administration in rats with an alternative nondrug reinforcer. *Psychopharmacology (Berl)* 2009; 203:13–22.
61. Domingos AI, Vaynshteyn J, Voss HU, *et al.* Leptin regulates the reward value of nutrient. *Nat Neurosci* 2011; 14:1562–1568.
62. Adamantidis AR, Tsai HC, Boutrel B, *et al.* Optogenetic interrogation of dopaminergic modulation of the multiple phases of reward-seeking behavior. *J Neurosci* 2011; 31:10829–10835.
63. Christensen CJ, Silberberg A, Hursh SR, *et al.* Essential value of cocaine and food in rats: tests of the exponential model of demand. *Psychopharmacology (Berl)* 2008; 198:221–229.
64. Christensen CJ, Silberberg A, Hursh SR, *et al.* Demand for cocaine and food over time. *Pharmacol Biochem Behav* 2008; 91:209–216.
65. Koffarnus MN, Woods JH. Individual differences in discount rate are associated with demand for self-administered cocaine, but not sucrose. *Addict Biol* 2013; 18:8–18.
66. Barnea-Ygael N, Yadid G, Yaka R, *et al.* Cue-induced reinstatement of cocaine seeking in the rat 'conflict model': effect of prolonged home-cage confinement. *Psychopharmacology (Berl)* 2012; 219:875–883.
67. Johnson PM, Kenny PJ. Dopamine D2 receptors in addiction-like reward dysfunction and compulsive eating in obese rats. *Nat Neurosci* 2010; 13: 635–641.
68. Oswald KD, Murdaugh DL, King VL, Boggiano MM. Motivation for palatable food despite consequences in an animal model of binge eating. *Int J Eat Disord* 2011; 44:203–211.
69. Pelloux Y, Everitt BJ, Dickinson A. Compulsive drug seeking by rats under punishment: effects of drug taking history. *Psychopharmacology (Berl)* 2007; 194:127–137.
70. Latagliata EC, Patrono E, Puglisi-Allegra S, Ventura R. Food seeking in spite of harmful consequences is under prefrontal cortical noradrenergic control. *BMC Neurosci* 2010; 11:15.